

The Three Faces of Ω_m : Testing Gravity with Low and High Redshift SN Ia Surveys

Alexandra Abate^{*} and Ofer Lahav[†]

Department of Physics & Astronomy, University College London, Gower Street, London, WC1E 6BT, UK

27 June 2008

ABSTRACT

Peculiar velocities of galaxies hosting Type Ia supernovae generate a significant systematic effect in deriving the dark energy equation of state w , at level of a few percent. Here we illustrate how the peculiar velocity effect in SN Ia data can be turned from a “systematic” into a probe of cosmological parameters. We assume a flat Λ -Cold Dark Matter model ($w = -1$) and use low and high redshift SN Ia data to derive simultaneously three distinct estimates of the matter density Ω_m which appear in the problem: from the geometry, from the dynamics and from the shape of the matter power spectrum. We find that each of the three Ω_m ’s agree with the canonical value $\Omega_m = 0.25$ to within 1σ , for reasonably assumed fluctuation amplitude and Hubble parameter. This is consistent with the standard cosmological scenario for both the geometry and the growth of structure. For fixed $\Omega_m = 0.25$ for all three Ω_m ’s, we constrain $\gamma = 0.72 \pm 0.21$ in the growth factor $\Omega_m(z)^\gamma$, so we cannot currently distinguish between standard Einstein gravity and predictions from some modified gravity models. Future surveys of thousands of SN Ia, or inclusion of peculiar velocity data, could significantly improve the above tests.

Key words: large-scale structure of universe – cosmological parameters – surveys – galaxies: kinematics and dynamics

1 INTRODUCTION

The observed present acceleration of the universe was first confirmed a decade ago by two separate groups using Type Ia supernovae (SN Ia, Perlmutter et al. 1999; Riess et al. 1998). SN Ia are one of a number of probes needed to obtain tighter constraints on dark energy equation of state, including any possible time evolution. This will require surveys of thousands of supernovae out to high redshifts to accurately measure their luminosity distances from which parameters describing the dark energy can be inferred. To achieve the desired constraints on dark energy, in particular a few percent constraint on the equation of state parameter w , the supernovae will have to be accurately calibrated. It is therefore vital that this calibration is done accurately, and it is the low redshift supernovae which are vital to achieve this, for details see Aldering et al. (2002). At low redshift the supernovae distances have little or no dependence on the cosmological parameters such as Ω_m , Ω_Λ and the dark energy equation of state w . They do however put a tight constraint on a combination of what is essentially the calibrated magnitude zeropoint (M) and the Hubble constant H_0 , whereas

for the high redshift supernovae there is a strong degeneracy between M , H_0 and the cosmological parameters of interest. Figure 1 illustrates the importance of the low redshift supernovae in anchoring the Hubble diagram. It shows the gold sample from Riess et al. (2007) constraints on Ω_m and w with and without supernovae with redshifts less than 0.1. One can see that without the low redshift supernovae (blue/light contours, using 146 SN Ia) the constraints blow up significantly compared to the full gold sample (red/dark contours, using 182 SN Ia). There are several sources of systematic error which affect the calibration of the zeropoint, for example dust extinction, luminosity evolution, weak lensing, and Malmquist bias (see Kim et al. 2004, for more details). This type of error is not decreased by having a large number of supernovae and will necessarily come to dominate the error budget. The systematic errors mentioned above have long been discussed in the literature and are not considered in this Letter. There is a source of error which is unique in the fact that it affects only the low redshift “calibrating” supernovae, their peculiar motions relative to the Hubble flow.

Previous authors have set about using SN Ia to quantify the degradation of dark energy errors due to peculiar motions or use them to trace the peculiar velocity field itself in a variety of ways. Three distinct approaches

^{*} E-mail: aabate@star.ucl.ac.uk

[†] E-mail: lahav@star.ucl.ac.uk

to this have been discussed recently in the literature. In Neill, Hudson, & Conley (2007) different flow models based on the IRAS PSCz survey (Branchini et al. 1999) were used to “correct” the luminosity distances by the known peculiar velocities before fitting them for the cosmological parameters of interest. They find the potential systematic error in w caused by ignoring peculiar velocities is of the order of 4 percent, i.e. quite significant.

Radburn-Smith, Lucey, & Hudson (2004) compared peculiar velocities from 98 local supernovae with the gravity field predicted from IRAS. In Haugbølle et al. (2007) an angular expansion of the radial velocity field was used to probe the local dipole and quadrupole of the velocity field at three different distances. They found that the dipole is consistent with galaxy surveys (e.g. Erdoğan et al. 2006) at the same Hubble flow depths.

The third and somewhat different method is utilised by Hui & Greene (2006); Cooray & Caldwell (2006); Gordon, Land & Slosar (2007) who take a covariance matrix approach. From the fluctuation in the luminosity distance induced by the peculiar motions, (see Hui & Greene 2006; Pyne & Birkinshaw 2004, 1996; Sasaki 1987, for derivations), a covariance matrix for the resulting errors in the luminosity distance (or similarly the apparent magnitude) can be calculated. The covariance matrix depends on cosmological parameters which describe the growth and distribution of structure. In addition to the peculiar velocity effect this is due to gravitational lensing effect, which is important for redshifts larger than 1, and we shall ignore it in this Letter.

Cooray & Caldwell (2006) found that peculiar velocities of the low redshift supernovae may prevent measurement of w to better than 10 percent, and diminish the resolution of the time derivative of w projected for planned surveys. Gordon, Land & Slosar (2007) used the covariance matrix approach on current data, showing the changing constraints on σ_8 , Ω_m and w depending on the exact redshift range of the SN Ia sample and whether the full covariance was included or not. They also apply the analysis to forecasting constraints for future surveys.

Here we unify the analysis of SN Ia data to study simultaneously fits for the expansion of the universe and the growth of structure. There is plenty of discussion on the possibility that the accelerated expansion of the universe is caused by a modification of general relativity on large scales (e.g. Durrer & Maartens 2008; Huterer & Linder 2007, and references therein). By measuring the growth of structure, which directly effects the observed peculiar velocity field, information is gained to differentiate between the two scenarios.

The rest of the Letter is organised as follows. In Section 2 we describe the SN Ia sample used in this Letter. In Section 3 we describe the theory underlying SN Ia analysis in cosmology and the effect of the velocity field.

2 DATA

We analyse both nearby supernovae ($z \leq 0.12$) from Jha, Riess, & Kirshner (2007) and high redshift supernovae ($z \leq 0.176$) from a sample compiled by Davis et al. (2007) which includes data from Riess et al. (2007) and Wood-Vasey et al. (2007). Davis et al. (2007) combined




Figure 1. Assuming a flat universe with a constant equation of state w , these are 1 and 2 σ likelihood contours showing the constraints in the Ω_m - w plane for the Riess et al 2007 supernova data. The red/dark contours are when using the full gold sample (182 SN Ia), the blue/light contours are when using the gold sample but removing all supernovae with redshifts less than 0.1 (146 SN Ia)




Figure 2. The 1 and 2 σ contours on Ω_m^{dyn} and the parameter γ from using all 271 low and high redshift SNe. Values of other parameters include $n_s = 1$ and $h = 0.7$, $\sigma_v = 300\text{kms}^{-1}$, $\sigma_m = 0.1$ and the other two Ω_m 's=0.25.

the data from the two samples by normalising to the low redshift supernovae they had in common. Following Jha, Riess, & Kirshner (2007) 9 supernovae are excluded from the low redshift set, those that are unsuitable due to bad lightcurve fits. This includes supernovae with their first observation more than 20 days after maximum light, those that are hosted in galaxies with excessive extinction ($A_V^0 > 2.0$ mag) and one outlier (SN1999e), which appears to have an extremely large peculiar velocity. This leaves 124 supernovae from the Jha, Riess, & Kirshner (2007) data set in the redshift range $z \in [0.0023, 0.12]$, and median redshift $\bar{z} = 0.017$. The overlapping SN Ia in the two data sets were used to estimate a small normalising offset to the magnitudes from the Davis et al. (2007) data set (the extra magnitude error is negligibly small). The same procedure was used by Davis et al. (2007) in normalising the two high redshift data sets. After eliminating duplicated SN Ia, our combined data set has 271 SNe with $z \in [0.0023, 1.76]$, and $\bar{z} = 0.29$.

3 METHODOLOGY

We describe here how we utilise the SN Ia dataset described in Section 2 to estimate cosmological parameters by including the peculiar velocity covariance.

3.1 Covariance matrix approach

The luminosity distance d_L is defined as

$$\mathcal{F} = \frac{\mathcal{L}}{4\pi d_L^2} \quad (1)$$

where $\mathcal{F}$ is the observed flux of the supernova and $\mathcal{L}$ is its intrinsic luminosity. The apparent magnitude m of a supernova at redshift z depends on the luminosity distance as follows

$$m(z) = 5 \log_{10} D_L(z) - 5 \log_{10}(H_0) + M + 25 \quad (2)$$

where M is the magnitude zeropoint, and D_L is defined without the Hubble constant as $D_L = H_0 d_l$ in kms^{-1} . The equation above ignores the additional terms which involve applying dust corrections, K corrections etc. For a flat universe containing a matter component and a dark energy component with a constant equation of state, the luminosity distance can be written as

$$D_L(z) = c(1+z) \int_0^z \frac{dz'}{E(z')} \quad (3)$$

$$E(z') = (\Omega_m(1+z')^3 + (1-\Omega_m)(1+z')^{-3(1+w)})^{\frac{1}{2}} \quad (4)$$

If the universe was truly homogeneous and isotropic (FRW) this would be the end of the story, the observed D_L would be described accurately by Eq. 3. However peculiar velocities have the effect of perturbing the luminosity distance

$$\frac{\delta D_L}{D_L} = \frac{v_r}{c} \left(1 - \frac{c(1+z)^2}{H(z)d_L(z)} \right) \quad (5)$$

where v_r is the radial peculiar velocity of the supernova and $H(z)$ is the Hubble parameter. See Hui & Greene (2006); Bonvin, Durrer, & Gasparini (2006); Pyne & Birkinshaw (2004); Sasaki (1987) for a derivation. We emphasize that in the right hand side of the above equation d_L is for an *unperturbed* FRW universe, derived at a *perturbed* redshift z .

Therefore the covariance of the perturbation in D_L , $\delta D_L/D_L$ for a pair i, j is given by

$$C_{ij}^L = \frac{\langle v_{ri} v_{rj} \rangle}{c^2} \left(1 - \frac{c(1+z)^2}{H(z)d_L(z)} \right)_i \left(1 - \frac{c(1+z)^2}{H(z)d_L(z)} \right)_j \quad (6)$$

and

$$\langle v_{ri} v_{rj} \rangle = \xi_{ij} = \cos \theta_i \cos \theta_j \Psi_{||}(r) + \sin \theta_i \sin \theta_j \Psi_{\perp}(r) \quad (7)$$

is the linear theory radial peculiar velocity correlation function, (Gorski 1988; Groth, Juszkiewicz, & Ostriker 1989). The angles in Eq. 7 are defined by $\cos \theta_X = \hat{\mathbf{r}}_X \cdot \hat{\mathbf{r}}$ and the diagonal elements ξ_{ii} are given by Eq. 9 below. The $\Psi_{||}(r)$ and $\Psi_{\perp}(r)$ can be calculated from the matter power spectrum using linear theory

$$\Psi_{||,\perp}(r) = D'(z_i)D'(z_j) \int \frac{P(k)}{2\pi^2} B_{||,\perp}(kr) dk \quad (8)$$




Figure 3. Constraints on Ω_m^{ps} and Ω_m^{dyn} from using all 271 low and high redshift SNe. In both panels Ω_m^{geom} has been fixed at 0.25, σ_v at 300 km s^{-1} and σ_m at 0.1. In the top panel n_s has been fixed at 1.0, and the red (dashed) contours, black (dark) contours and blue (light) contours are the results when $h = 0.6$, $h = 0.7$ and $h = 0.8$ respectively. In the bottom panel h has been fixed at 0.7, and the red (dashed) contours, black (dark) contours and blue (light) contours are the results when $n_s = 0.9$, $n_s = 1$ and $n_s = 1.1$ respectively. The black (dark) contours in both panels are exactly the same. The grey line indicates where $\Omega_m^{ps} = \Omega_m^{dyn}$.

where $P(k)$ is the matter power spectrum, $B_{\perp} = j_0'(x)/x$ and $B_{||} = j_0''(x)$ and j_0', j_0'' are the first and second derivative of the zeroth order spherical Bessel functions respectively and $D'(z)$ is the derivative of the growth function at redshift z . $D'(z)$ is a function of the Hubble parameter and Ω_m . See Section 3.3 for further discussion. The auto correlation is given by

$$\xi_{ii} = \frac{1}{3} D'^2(z_i) \int \frac{P(k)}{2\pi^2} dk. \quad (9)$$

We can therefore calculate C_{ij}^L for a pair of supernovae at z_i and z_j respectively given a set of cosmological parameters.

In the above equations the growth factor is calculated exactly numerically. More insight to the dependence on Ω_m is given by the commonly used approximation for the growth factor $f = d \ln \delta / d \ln a \approx \Omega_m(z)^\gamma$, where $\gamma \approx 0.6$ (Peebles 1980), with little dependence on the cosmological constant (Lahav et al. 1991), and a slight dependence on w (Wang & Steinhardt 1998). Recent refined calculations predict $\gamma = 0.55$ for the concordance model, and $\gamma = 0.69$ (Linder & Cahn 2007) for a particular modified gravity model, DGP braneworld gravity (Dvali, Gabadadze, & Porrati 2000), though this is just an example of many possible modified gravity models. Below we shall constrain γ from the SN Ia data.

3.2 Likelihood analysis

To find the set of cosmological parameters $\Theta_{\max} = [\theta_1 \dots \theta_N]$ that best fit the data we find the set that maximise the likelihood function. Assuming that the data and the observational errors are Gaussian random fields the likelihood function can be written as

$$\mathcal{L} = \frac{1}{\sqrt{(2\pi)^N |\Sigma|}} \exp \left(-\frac{1}{2} \sum_{i,j} D_i (\Sigma^{-1})_{ij} D_j \right). \quad (10)$$

where D_i is defined as $D_i = (D_L^{obs} - D_L(z)) / D_L(z)$ and Σ is the covariance matrix including the observational noise. Following Gordon, Land & Slosar (2007) we write this as

$$\Sigma_{ij} = C_{ij}^L + \sigma_i^2 \delta_{ij} \quad (11)$$

where σ_i is the standard uncorrelated error given by

$$\sigma_i^2 = \left(\frac{\ln(10)}{5} \right)^2 (\sigma_m^2 + (\mu_i^{err})^2) + \left(1 - \frac{c(1+z)^2}{H(z)d_L(z)} \right)^2 \frac{\sigma_v^2}{c^2} \quad (12)$$

where σ_v is often set to 300 km s^{-1} and is included to account for nonlinear contributions to ξ_{ij} (which is derived only in linear theory, Silberman et al. 2001), and the velocity of the SN within the host galaxy. Here σ_m is the intrinsic magnitude scatter and μ^{err} is the error from the light curve fitting.

3.3 The Three Faces of Ω_m

From the equations in Section 3.1 it can easily be seen that C_{ij}^L is a function of Ω_m through

(i) Ω_m^{geom} : the geometry of the universe, from $H(z)$ and also $d_L(z)$ in Eq. 5. For a flat universe $\Omega_m^{\text{geom}} = 1 - \Omega_\Lambda$, with no dependence on other cosmological parameters. Ω_m^{geom} is most strongly constrained however by the high redshift SNe through Eqs. 3 and 4.

(ii) Ω_m^{dyn} : the growth of structure in the universe, from $D'(z)$ in Eqs. 8 and 9 or the equivalent growth factor parameterization $f = [\Omega_m^{\text{dyn}}(z)]^\gamma$. We note a strong degeneracy through the product $\sigma_8 \Omega_m(z)^\gamma$ where

$$\Omega_m(z) = \frac{\Omega_m(z=0)(1+z)^3}{\Omega_m(z=0)(1+z)^3 + 1 - \Omega_m(z=0)} \quad (13)$$

for a flat universe, our results can be scaled accordingly.

(iii) Ω_m^{ps} : the matter power spectrum $P(k)$ in Eqs. 8 and 9. It is well known that the shape of the power spectrum depend on the product $\Gamma = \Omega_m^{\text{ps}} h$, with some degeneracy with e.g. the spectral index n_s , σ_8 and baryon and neutrino mass densities Ω_b and Ω_ν .

Please note that Eq.'s 8 and 9 contain all of the low redshift Ω_m terms. If the Λ CDM model of the universe is correct then when varying each of these “faces” of Ω_m separately the results should be consistent with each other. If not this suggests that the Λ CDM model is inconsistent and the data may favour a model which changes the theory of general relativity on large scales or other dark energy models.

Table 1. Results for each “face” of Ω_m under different parameter combinations. Because we do not marginalise the errors are small, but it is still useful to look at the relative errors for the three Ω_m 's. Columns A, C and D set the other two $\Omega_m = 0.25$, whereas column B marginalises over them. In column A and B the “nuisance” parameters are $\sigma_v = 300 \text{ km s}^{-1}$ and $\sigma_m = 0.1$. Column C assumes the “nuisance” parameters are small ($\sigma_v = 200 \text{ km s}^{-1}$ and $\sigma_m = 0.08$). Column D assumes the “nuisance” parameters are large ($\sigma_v = 400 \text{ km s}^{-1}$ and $\sigma_m = 0.12$). Each column has set the power spectrum parameters $[h \ n_s \ \sigma_8] = [0.7 \ 1.0 \ 0.8]$.

	A	B	C	D
Ω_m^{geom}	$0.25^{+0.01}_{-0.01}$	$0.25^{+0.01}_{-0.01}$	$0.25^{+0.01}_{-0.01}$	$0.24^{+0.01}_{-0.01}$
Ω_m^{dyn}	$0.25^{+0.14}_{-0.14}$	$0.18^{+0.11}_{-0.11}$	$0.33^{+0.14}_{-0.14}$	$0.10^{+0.10}_{-0.14}$
Ω_m^{ps}	$0.43^{+0.46}_{-0.26}$	$0.38^{+0.45}_{-0.25}$	$0.33^{+0.43}_{-0.23}$	$0.48^{+0.52}_{-0.26}$

4 RESULTS

In the following analysis we assume a flat Λ CDM universe with a dark energy equation of state $w = -1$. To gain an insight for the effect of varying the other cosmological parameters, namely H_0 , n_s , and the “nuisance” parameters σ_m and σ_v we do not marginalise over them but present the results at some choice values for these parameters. The effect of marginalising over σ_v and σ_m degrades the error on Ω_m^{geom} by about 10 percent, the error on Ω_m^{ps} changes negligibly and the error on Ω_m^{dyn} degrades by about 30 percent. The larger degradation of the error on Ω_m^{dyn} is because of its strong degeneracy with σ_v . Other parameters are fixed as follows; $\Omega_b = 0.04$, $\Omega_\nu = 0$ and $\sigma_8 = 0.8$. For clarity most contours have only the 1σ confidence level, and where appropriate the assumed values of other parameters are stated.

Table 1 shows a set of results for each “face” of Ω_m under different parameter combinations. This table shows that the different Ω_m 's are consistent with the canonical value of 0.25 to within 1σ . One can see the degeneracy direction of the errors (σ_v , σ_m) and Ω_m^{dyn} , discussed below. See table caption for an explanation of the columns.

Figure 2 shows the constraints on Ω_m^{dyn} and the parameter γ , described at the end of Section 3.1. The contours shown are the 1 and 2σ . For $\Omega_m^{\text{dyn}} = 0.25$ the 68 percent confidence constraint on γ is $\gamma = 0.72 \pm 0.21$ consistent with both Einstein gravity ($\gamma = 0.55$) and DGP gravity ($\gamma = 0.69$).

Figure 3 shows the effect of h and n_s on the Ω_m^{ps} - Ω_m^{dyn} contours. As expected there is only a strong influence on Ω_m^{ps} . One can also see the degeneracy direction of h and n_s in the power spectrum, shown by the indistinguishable differences between the red (dashed) contours in both panels. The positions of the red (dashed) contours show that decreasing h by roughly 10 percent is equivalent to increasing n_s also by roughly 10 percent. This is also shown by the blue (light) contours. The contours for Ω_m^{ps} - Ω_m^{geom} and Ω_m^{geom} - Ω_m^{dyn} are not shown here since the same combinations of h and n_s as plotted in Figure 3 have a nearly negligible effect on the contour positions. Under different permutations of $h = 0.6, 0.7, 0.8$ and $n_s = 0.9, 1.0, 1.1$ the contours only shift in either the Ω_m^{ps} or Ω_m^{dyn} direction and even then there is only a maximum shift of $\Delta\Omega_m$ of 0.05.




Figure 4. Constraints on all the Ω_m pairs from using all 271 low and high redshift SNe. From the left the first panel shows Ω_m^{ps} - Ω_m^{dyn} , the second panel Ω_m^{ps} - Ω_m^{geom} , third panel Ω_m^{geom} - Ω_m^{dyn} and the fourth panel Ω_m^{dyn} - σ_v . The black (dark), red (dashed) and blue (light) contours in the first 3 panels correspond to σ_v equal to 250km s^{-1} , 300km s^{-1} and 350km s^{-1} respectively. Each time $n_s = 1$, $h = 0.7$, $\sigma_m = 0.1$, $\sigma_8 = 0.8$ and in the first 3 panels the third Ω_m is kept fixed at 0.25 and just the $1\text{-}\sigma$ contour is shown. In the fourth panel $n_s = 1$, $h = 0.7$, $\sigma_m = 0.1$, $\Omega_m^{ps} = \Omega_m^{geom} = 0.25$ and both the 1 and $2\text{-}\sigma$ contours are shown. The grey line indicates where $\Omega_m^x = \Omega_m^y$.

Finally Figure 4 shows the effect of σ_v on all the contour pairs. It has the largest effect on Ω_m^{dyn} and very little effect on Ω_m^{geom} . This is because the diagonal elements of C_{ij}^L (see Eq. 6) are approximately proportional to $(\Omega_m^{dyn})^{2\gamma}$ and to σ_v^2 . Therefore the best-fit Ω_m^{dyn} decreases as σ_v increases.

5 CONCLUSIONS

We have presented in this paper a unified approach for probing both the expansion of the universe and the growth of structure with SN Ia data and to test the consistency of the Λ CDM model. We utilised the SNIa data to derive three distinct estimates of the matter density Ω_m which appear in the problem: from the geometry, from the dynamics and from the shape of the matter power spectrum. We found that each of them agrees with canonical value $\Omega_m = 0.25$ to within 1σ . We note we are restricting our discussion to Λ CDM, if we allow w to vary our constraints on Ω_m^{geom} will weaken. We also constrained γ in the growth factor $\Omega_m(z)^\gamma$ and found for $\Omega_m = 0.25$, $\gamma = 0.72 \pm 0.21$. This value of γ is consistent with both concordance and some proposed modified gravity models.

Current and future SN Ia surveys such as SN Factory, GAIA and Skymapper (for low redshift), SDSS-II (for intermediate redshift) and DES, Pan-STARRS, LSST, DUNE and SNAP (for high redshift) will generate samples of thousands of SN Ia (e.g. Albrecht et al. 2006; Peacock et al. 2006, for overviews). Large samples of low redshift SNe will greatly improve our constraints on Ω_m^{ps} and Ω_m^{dyn} and the high redshift SNe on Ω_m^{geom} . Utilising *galaxy* peculiar velocity data (using $D_n - \sigma$ and Tully-Fisher distance indicators) will also provide improvement on the Ω_m^{ps} and Ω_m^{dyn} constraints. Our approach can also be generalised for a range of other cosmological parameters and exotic models of dark energy and gravity.

6 ACKNOWLEDGMENTS

We are grateful to Kate Land for providing us with information about SN Ia data sets, and to Sarah Bridle, Josh Frieman, Chris Gordon, Saurabh Jha, John Marriner and Jochen Weller for useful conversations. AA acknowledges the receipt of a STFC studentship and OL acknowledges a Royal Society Wolfson Research Merit Award. Both AA and

OL thank Fermilab and the Kavli Institute for Cosmological Physics at Chicago University for their hospitality. We thank the anonymous referee for their detailed comments.

REFERENCES

- Albrecht A., et al., 2006, *astro*, arXiv:astro-ph/0609591
- Aldering G., et al., 2002, *SPIE*, 4836, 61
- Bonvin C., Durrer R., Gasparini M. A., 2006, *PhRvD*, 73, 023523
- Branchini E., et al., 1999, *MNRAS*, 308, 1
- Bridle S. L., Crittenden R., Melchiorri A., Hobson M. P., Kneissl R., Lasenby A. N., 2002, *MNRAS*, 335, 1193
- Cooray A., Caldwell R. R., 2006, *PhRvD*, 73, 103002
- Davis T. M., et al., 2007, *ApJ*, 666, 716
- Durrer R., Maartens R., 2008, *GRGr*, 40, 301
- Dvali G., Gabadadze G., Porrati M., 2000, *PhLB*, 485, 208
- Erdođdu P., et al., 2006, *MNRAS*, 368, 1515
- Gordon C., Land K., Slosar A., 2007, *PhRvL*, 99, 081301
- Gorski K., 1988, *ApJ*, 332, L7
- Groth E. J., Juskiewicz R., Ostriker J. P., 1989, *ApJ*, 346, 558
- Haugbølle T., Hannestad S., Thomsen B., Fynbo J., Sollerman J., Jha S., 2007, *ApJ*, 661, 650
- Hui L., Greene P. B., 2006, *PhRvD*, 73, 123526
- Huterer D., Linder E. V., 2007, *PhRvD*, 75, 023519
- Kim A. G., Linder E. V., Miquel R., Mostek N., 2004, *MNRAS*, 347, 909
- Jelinsky P., SNAP Collaboration, 2006, *AAS*, 209, #98.09
- Jha S., Riess A. G., Kirshner R. P., 2007, *ApJ*, 659, 122
- Kolb E. W., Matarrese S., Notari A., Riotto A., 2005, *PhRvD*, 71, 023524
- Lahav O., Lilje P. B., Primack J. R., Rees M. J., 1991, *MNRAS*, 251, 128
- Linder E. V., Cahn R. N., 2007, *APh*, 28, 481
- Neill J. D., Hudson M. J., Conley A., 2007, *ApJ*, 661, L123
- Peacock J. A., Schneider P., Efstathiou G., Ellis J. R., Leibundgut B., Lilly S. J., Mellier Y., 2006, *astro*, arXiv:astro-ph/061090
- Peebles P. J. E., 1980, *The Large Scale Structure of the Universe*, Princeton University Press
- Peel A., Knox L., 2003, *NuPhS*, 124, 83
- Perlmutter S., et al., 1999, *ApJ*, 517, 565
- Pyne T., Birkinshaw M., 1996, *ApJ*, 458, 46
- Pyne T., Birkinshaw M., 2004, *MNRAS*, 348, 581

- Radburn-Smith D. J., Lucey J. R., Hudson M. J., 2004, MNRAS, 355, 1378
- Riess A. G., et al., 1998, AJ, 116, 1009
- Riess A. G., et al., 2007, ApJ, 659, 98
- Sasaki M., 1987, MNRAS, 228, 653
- Silberman L., Dekel A., Eldar A., Zehavi I., 2001, ApJ, 557, 102
- Sugiura N., Sugiyama N., Sasaki M., 1999, PThPh, 101, 903
- Wang L., Steinhardt P. J., 1998, ApJ, 508, 483
- Wood-Vasey W. M., et al., 2002, AAS, 34, 1205
- Wood-Vasey W. M., et al., 2007, ApJ, 666, 694